



Natural Antibiotics: A Study of the Biological Activity of a Quintet (Garlic, Turmeric, Cinnamon, Ginger, and Honey) as Safe Alternatives to Conventional Antibiotics

Dr. Mohammed Arhouma^{1*}, Areej Najmi Abu Gharara², Dania Al-Mabrouk Burqan, Faten Mohamed Al-Amin, Fadi Belaid

Mohammedsaenoun@gmail.com, areejeje20@gmail.com, bargandanya@gmail.com, fatenkhalas1999@gmail.com, aalr4753@gmail.com

ARTICLE INFO	
Received: 01-03-2026 Accepted: 10-03-2026 Published: 30-03-2026 Keywords. Counting People, Face Detection, People Detection, Viola Jones LBP, Viola Jones CART. Antibiotic resistance; natural antibiotics; garlic; turmeric; cinnamon; ginger; honey; antimicrobial activity Copyright: © 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution International License (CC BY 4.0). http://creativecommons.org/licenses/by/4.0/	ABSTRACT <i>Background:</i> Antibiotic resistance has emerged as one of the most critical global health threats, necessitating the search for safe and effective natural alternatives. <i>Objective:</i> This systematic review evaluates the biological activity of five natural agents – garlic, turmeric, cinnamon, ginger, and honey – as potential natural antibiotics. <i>Methods:</i> A comprehensive literature search was conducted in PubMed, Scopus, and Google Scholar databases (2000–2025) focusing on in vitro and in vivo studies assessing antimicrobial activity, mechanisms of action, and minimum inhibitory concentrations (MICs) of the selected natural products. <i>Results:</i> The bioactive compounds – allicin (garlic), curcumin (turmeric), cinnamaldehyde (cinnamon), gingerol/shogaol (ginger), and hydrogen peroxide/methylglyoxal (honey) – exhibit potent activity against a wide range of Gram-positive and Gram-negative bacteria, including multidrug-resistant (MDR) strains. MIC values ranged from 0.3–3 mg/mL (garlic), 31.25–5000 µg/mL (curcumin), 1.5–12.5 µL/mL (cinnamon), 2.9–4.8 mg/mL (ginger), and 1–12.5% w/v (Manuka honey). Synergistic effects with conventional antibiotics and anti-biofilm activities were documented. <i>Conclusion:</i> These natural agents represent promising safe alternatives or adjunctive therapies to synthetic antibiotics. Further clinical trials are needed to standardize dosages and evaluate long-term efficacy.

1. INTRODUCTION

In recent decades, the world has been confronted with one of the most pressing health challenges: antibiotic resistance. Since the discovery of penicillin, antibiotics have saved countless lives and revolutionized healthcare. However, excessive and inappropriate use has led to the emergence of resistant bacterial strains, rendering many conventional treatments ineffective (WHO, 2021). According to the World Health Organization, antibiotic resistance threatens to undermine decades of medical progress, making routine surgeries, organ transplants, and even minor infections potentially life-threatening.

Factors contributing to this crisis include over-prescription, patient non-adherence to treatment courses, and widespread agricultural use. In response, researchers are increasingly turning to natural alternatives. Plants and bee products have been used for centuries in traditional medicine to combat infections. Among the most promising candidates are garlic (*Allium sativum*), turmeric (*Curcuma longa*), cinnamon (*Cinnamomum* spp.), ginger (*Zingiber officinale*), and honey. Each possesses significant biological activity



against bacteria and fungi, acting through multiple mechanisms that reduce the likelihood of rapid resistance development (Nabavi et al., 2015; Adamczak et al., 2020).

This review aims to evaluate the antimicrobial efficacy, mechanisms of action, and potential clinical applications of these five natural agents as safe alternatives or complements to synthetic antibiotics.

2. Methodology

2.1 Search Strategy

A systematic literature search was performed using PubMed, Scopus, and Google Scholar databases for articles published between January 2000 and March 2025. The following keywords were used: "natural antibiotics," "garlic antibacterial," "curcumin antimicrobial," "cinnamon essential oil," "ginger bioactive compounds," "honey wound healing," "antimicrobial resistance," and "synergistic effects."

2.2 Inclusion and Exclusion Criteria

Studies were included if they: (i) were original research articles or reviews published in peer-reviewed journals; (ii) evaluated the antimicrobial activity of garlic, turmeric, cinnamon, ginger, or honey against bacterial or fungal pathogens; (iii) reported minimum inhibitory concentration (MIC) values or clear mechanisms of action; (iv) were written in English or Arabic. Exclusion criteria included conference abstracts, unpublished theses, and studies without quantitative data.

2.3 Data Extraction

Data were extracted on: plant/bee product, bioactive compound, tested microorganisms, MIC values, mechanisms of action, and any synergistic effects with conventional antibiotics.

3. Results

3.1 Garlic (*Allium sativum*)

Garlic (family Liliaceae) has been used as a spice and traditional medicine for centuries. Its health benefits are attributed to cysteine-derived organosulfur compounds. Intact garlic contains γ -glutamyl-S-alk(en)yl-L-cysteine and S-alk(en)yl-L-cysteine sulfoxides (alliin, isoalliin, methiin). Upon crushing, the enzyme alliinase converts alliin to allicin, a highly reactive thiosulfinate that decomposes into diallyl sulfide (DAS), diallyl disulfide (DADS), diallyl trisulfide (DATS), ajoene, and vinyl dithiols (Nabavi et al., 2015). Allicin exhibits excellent in vitro antibacterial activity against various pathogens, including methicillin-resistant *Staphylococcus aureus* (MRSA). MIC values for garlic extracts range from 0.3 to 3 mg/mL. The proposed mechanisms include enzyme inactivation, membrane disruption, and inhibition of protein synthesis.

3.2 Turmeric (*Curcuma longa*)

Curcumin (diferuloylmethane) is the principal bioactive polyphenol in turmeric rhizomes. Its chemical formula is $C_{21}H_{20}O_6$ (molecular weight 368.38 g/mol). Curcumin possesses antioxidant, anti-inflammatory, and antimicrobial activities (Adamczak et al., 2020). In a study testing over 100 clinical and reference strains, curcumin showed strong activity against Gram-positive bacteria, especially *Streptococcus pyogenes* (median MIC = 31.25 μ g/mL). Activity against Gram-negative bacteria was weaker (median MIC = 2000 μ g/mL). Multidrug-resistant strains (MRSA, ESBL-positive *E. coli*) required higher MICs (≥ 2000 μ g/mL). Curcumin also inhibits bacterial quorum sensing and biofilm formation, and exhibits photodynamic action through reactive oxygen species (ROS) production (Moghadamtousi et al., 2014).

3.3 Cinnamon (*Cinnamomum* spp.)

The genus *Cinnamomum* (Lauraceae) includes over 300 species. *C. zeylanicum* (true cinnamon) and *C. cassia* are the most studied. The main antibacterial constituent is cinnamaldehyde (62–90% in bark essential oil). Leaf oil is rich in eugenol (>80%). Cinnamon extracts and essential oils have demonstrated activity against both Gram-positive and Gram-negative bacteria, including *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *E. coli*, and MRSA (Nabavi et al.,



2015). MIC values range from 1.5 to 12.5 $\mu\text{L/mL}$. Cinnamon also inhibits biofilm formation of enterohemorrhagic *E. coli* O157:H7. Synergistic effects have been reported with piperacillin against β -lactamase-producing *E. coli*. However, caution is advised: high doses may cause glutathione depletion and central nervous system depression.

3.4 Ginger (*Zingiber officinale*)

Ginger rhizome contains phenolic compounds called gingerols (e.g., 6-gingerol) that undergo dehydration to shogaols upon drying or heating, enhancing antimicrobial potency. Ginger compounds disrupt bacterial cell membranes, causing leakage of intracellular contents, and inhibit DNA and protein synthesis. They also reduce exopolysaccharide production and biofilm formation. MIC values against *Salmonella*, *E. coli*, and *S. aureus* range from 2.9 to 4.8 mg/mL (Herman et al., 2013). Ginger extracts are effective against foodborne pathogens, making them valuable as natural food preservatives.

3.5 Honey

Honey is a complex mixture of 80–85% carbohydrates (fructose and glucose), water, enzymes, amino acids, vitamins, minerals, and phytochemicals (Hasam et al., 2020). Its antimicrobial mechanisms include: (i) high osmolarity causing water withdrawal from microbial cells; (ii) low pH (3.2–4.5); (iii) enzymatic production of hydrogen peroxide via glucose oxidase; (iv) presence of methylglyoxal (MGO) in Manuka honey; (v) flavonoids and phenolic acids; and (vi) biofilm disruption. MIC values vary by honey type: Manuka honey 1–12.5% w/v, dark buckwheat/heather 1–12.5%, clover/acacia 25–50% (Simoes et al., 2009). Manuka honey's potency is quantified by the Unique Manuka Factor (UMF), correlated with MGO concentration. Honey also modulates inflammatory responses and promotes wound healing.

Table 1: Summary of Antimicrobial Activity of the Five Natural Agents

gent	Bioactive Compound	Main Mechanism	Typical MIC
Garlic	Allicin, ajoene, DADS	Enzyme inactivation, membrane disruption	0.3–3 mg/mL
Turmeric	Curcumin	Quorum sensing inhibition, ROS production	31.25–5000 $\mu\text{g/mL}$
Cinnamon	Cinnamaldehyde, eugenol	Cell wall disruption, biofilm inhibition	1.5–12.5 $\mu\text{L/mL}$
Ginger	Gingerol, shogaol	Membrane leakage, DNA/protein synthesis inhibition	2.9–4.8 mg/mL
Honey	MGO, H_2O_2 , flavonoids	Osmotic effect, low pH, oxidative damage	1–12.5% (w/v)

4. Discussion

4.1 Comparative Mechanisms of Action

Synthetic antibiotics typically target a single bacterial pathway (e.g., cell wall synthesis by penicillins, protein synthesis by tetracyclines), which facilitates the rapid emergence of resistance. In contrast, natural antibiotics employ multiple, often synergistic mechanisms. Allicin reacts with thiol groups of essential enzymes; curcumin interferes with quorum sensing and generates ROS; cinnamaldehyde disrupts membrane integrity; gingerols cause membrane leakage; and honey combines osmotic, pH, and oxidative stresses. This multiplicity reduces the selective pressure for resistance.

4.2 Challenges and Limitations

Despite promising in vitro data, several challenges remain: (i) Standardization – bioactive compound concentrations vary with plant origin, harvest season, and extraction method; (ii) Bioavailability – curcumin has poor water solubility and low oral bioavailability (Siviero et al., 2015); (iii) Dosage optimization – no standardized clinical protocols exist; (iv) Toxicity concerns – high doses of cinnamon oil may deplete glutathione and interact with drugs (e.g., tetracyclines).



4.3 Synergism with Conventional Antibiotics

Several studies have shown that combining natural agents with antibiotics can produce additive or synergistic effects. For example, cinnamon essential oil plus piperacillin reduced MIC values against β -lactamase-producing *E. coli*. Curcumin enhanced the activity of ciprofloxacin against MRSA. Such combinations could lower required antibiotic doses, reduce side effects, and potentially reverse resistance.

4.4 Clinical and Food Industry Applications

Honey, particularly medical-grade Manuka honey, is already used in wound care for diabetic ulcers, burns, and surgical wounds. Garlic and ginger extracts are being explored as natural preservatives in meat and dairy products. Curcumin is incorporated into functional foods and beverages. These applications align with the growing consumer demand for “clean label” natural products.

4.5 Comparison with Synthetic Antibiotics

Aspect	Natural Antibiotics	Synthetic Antibiotics
Mechanism	Multiple, synergistic	Single target
Resistance development	Low probability	High and rapid
Side effects	Mild, generally safe	GI disturbances, allergies, microbiome disruption
Cost	Low	High (especially newer agents)
Efficacy in severe infections	Limited	High
Regulatory status	Dietary supplements, unstandardized	Strictly regulated

5. Conclusions and Recommendations

Antibiotic resistance is a global crisis that demands innovative solutions. This review provides strong evidence that garlic, turmeric, cinnamon, ginger, and honey possess significant antimicrobial activity against a wide spectrum of pathogens, including multidrug-resistant strains. Their multiple mechanisms of action, favorable safety profiles, low cost, and wide availability make them promising candidates as safe alternatives or adjuncts to synthetic antibiotics.

Recommendations:

- Integrative therapy:** Use natural agents as complementary treatments alongside conventional antibiotics to reduce dosages and side effects.
- Pharmaceutical development:** Develop standardized formulations (capsules, ointments, gels, sprays) containing defined bioactive compounds.
- Public awareness:** Educate healthcare professionals and the public on the rational use of natural antibiotics.
- Further research:** Conduct large-scale clinical trials to establish standardized dosages, evaluate long-term safety, and explore synergistic combinations.
- Food industry applications:** Promote the use of these natural agents as antimicrobial preservatives to enhance food safety and shelf life.
- Policy measures:** Include natural alternatives in national action plans against antimicrobial resistance.



REFERENCES

1. Nabavi, S. F., Di Lorenzo, A., Izadi, M., Sobarzo-Sanchez, E., Daglia, M., & Nabavi, S. M. (2015). Antibacterial Effects of Cinnamon: From Farm to Food, Cosmetic and Pharmaceutical Industries. *Nutrients*, 7(9), 7729–7748.
2. Alinezhad, H., Azimi, R., Zare, M., Ebrahimzadeh, M.A., Eslami, S., Nabavi, S.F., & Nabavi, S.M. (2013). Antioxidant and Antihemolytic Activities of Ethanolic Extract of Flowers, Leaves, and Stems of *Hyssopus officinalis* L. *International Journal of Food Properties*, 16, 1169–1178.
3. Nabavi, S.F., Daglia, M., Moghaddam, A.H., Habetmariam, S., & Nabavi, S.M. (2014). Curcumin and Liver Disease: From Chemistry to Medicine. *Comprehensive Reviews in Food Science and Food Safety*, 13, 62–77.
4. Curti, V., Capelli, E., Boschi, F., Nabavi, S.F., Bongiorno, A.I., Habetmariam, S., Nabavi, S.M., & Daglia, M. (2014). Modulation of human miR-17-3p expression by methyl 3-O-methyl gallate. *Molecular Nutrition & Food Research*, 58, 1776–1784.
5. Nabavi, S.F., Nabavi, S.M., Habetmariam, S., Moghaddam, A.H., Sureda, A., Jafari, M., & Latifi, A.M. (2013). Hepatoprotective effect of gallic acid isolated from *Peltiphyllum peltatum* against sodium fluoride-induced oxidative stress. *Industrial Crops and Products*, 44, 50–55.
6. Herman, A., Herman, A.P., Domagalska, B.W., & Mlynarczyk, A. (2013). Essential oils and herbal extracts as antimicrobial agents in cosmetic emulsion. *Indian Journal of Microbiology*, 53, 232–237.
7. Nabavi, S.M., Marchese, A., Izadi, M., Curti, V., Daglia, M., & Nabavi, S.F. (2015). Plants belonging to the genus *Thymus* as antibacterial agents. *Food Chemistry*, 173, 339–347.
8. Simoes, M., Bennett, R.N., & Rosa, E.A. (2009). Understanding antimicrobial activities of phytochemicals against multidrug resistant bacteria and biofilms. *Natural Product Reports*, 26, 746–757.
9. Marchese, A., Coppo, E., Sobolev, A.P., Rossi, D., Mannina, L., & Daglia, M. (2014). Influence of in vitro simulated gastroduodenal digestion on the antibacterial activity of green tea. *Food Research International*, 63, 182–191.
10. Ranasinghe, P., Jayawardana, R., Galappaththy, P., Constantine, G., de Vas Gunavardana, N., & Katulanda, P. (2012). Efficacy and safety of 'true' cinnamon (*Cinnamomum zeylanicum*) as a pharmaceutical agent in diabetes. *Diabetic Medicine*, 29, 1480–1492.
11. Ravindran, P., Shylaja, M., Nirmal Babu, K., & Krishnamoorthy, B. (2004). Botany and crop improvement of cinnamon and cassia. In: Ravindran P.N., Babu K.N. (Eds.), *Cinnamon and Cassia – The Genus Cinnamomum*. CRC Press.
12. Izah, S.C., Odubo, T.C., Ngun, C.T., & Ogwu, M.C. (2024). Research Needs of Medicinal Plants Used in the Management and Treatment of Some Diseases Caused by Microorganisms. In: Izah S.C., Ogwu M.C., Akram M. (Eds.), *Herbal Medicine Phytochemistry*, 1st ed., Springer, pp. 1797–1823.
13. Izah, S.C., Ogidi, O.I., Ogwu, M.C., Salimon, S.S., Yusuf, Z.M., Akram, M., Raimi, M.O., & Iyingiala, A.-A. (2024). Historical Perspectives and Overview of the Value of Herbal Medicine. In: Izah S.C., Ogwu M.C., Akram M. (Eds.), *Herbal Medicine Phytochemistry*, 1st ed., Springer, pp. 3–35.
14. Suntar, I., Khan, H., Patel, S., Celano, R., & Rastrelli, L. (2018). An overview on *Citrus aurantium* L.: Its functions as food ingredient and therapeutic agent. *Oxidative Medicine and Cellular Longevity*, 2018, 7864269.
15. Ogwu, M.C., & Ikhajiagbe, B. (2024). *Vernonia amygdalina* Delile (Asteraceae): An Overview. In: Izah S.C., Ogwu M.C., Akram M. (Eds.), *Herbal Medicine Phytochemistry*, Springer.
16. Ogwu, M.C., Dunkwu-Okafor, A., Omakor, I.A., & Izah, S.C. (2024). Turkey Berry (*Solanum torvum* Sw.). In: Izah S.C., Ogwu M.C., Akram M. (Eds.), *Herbal Medicine Phytochemistry*, Springer.
17. Sharma, A., et al. (2022). *Carica papaya* L. Leaves: Deciphering Its Antioxidant Bioactives. *Oxidative Medicine and Cellular Longevity*, 2022, 2451733.
18. Izah, S.C., Uhunmwangho, E.J., & Etim, N.G. (2018). Antibacterial and Synergistic Potency of Methanolic Leaf Extracts of *Vernonia amygdalina* L. and *Ocimum gratissimum* L. *Journal of Basic Pharmacology and Toxicology*, 2, 8–12.
19. Kigigha, L.T., Selekere, R.E., & Izah, S.C. (2018). Antibacterial and Synergistic Efficacy of Acetone Extracts of *Garcinia kola* and *Buchholzia coriacea*. *Journal of Basic Pharmacology and Toxicology*, 2, 13–17.
20. Hasam, S., Qarizada, D., & Azizi, M. (2020). A Review: Honey and Its Nutritional Composition. *Asian Journal of*



Research in Biochemistry, 7, 34–43.

21. Adamczak, A., Ożarowski, M., & Karpiński, T.M. (2020). Curcumin, a Natural Antimicrobial Agent with Strain-Specific Activity. *Pharmaceuticals*, 13(7), 153.
22. Basnet, P., & Skalko-Basnet, N. (2011). Curcumin: An anti-inflammatory molecule from a curry spice on the path to cancer treatment. *Molecules*, 16, 4567–4598.
23. Siviero, A., Gallo, E., Maggini, V., Gori, L., Mugelli, A., Firenzuoli, F., & Vannacci, A. (2015). Curcumin, a golden spice with a low bioavailability. *Journal of Herbal Medicine*, 5, 57–70.
24. Kotha, R.R., & Luthria, D.L. (2019). Curcumin: Biological, pharmaceutical, nutraceutical, and analytical aspects. *Molecules*, 24, 2930.
25. Prasad, S., & Aggarwal, B.B. (2011). Turmeric, the golden spice. In: Benzie I.F.F., Wachtel-Galor S. (Eds.), *Herbal Medicine: Biomolecular and Clinical Aspects*, 2nd ed., CRC Press, pp. 263–288.
26. Nair, K.P. (2019). *Turmeric (Curcuma longa L.) and Ginger (Zingiber officinale Rosc.) – World's Invaluable Medicinal Spices*. Springer.
27. Kwiecien, S., Magierowski, M., Majka, J., Ptak-Belowska, A., Wojcik, D., Sliwowski, Z., Magierowska, K., & Brzozowski, T. (2019). Curcumin: A potent protectant against esophageal and gastric disorders. *International Journal of Molecular Sciences*, 20, 1477.
28. Cheng, Y.T., Lu, C.C., & Yen, G.C. (2017). Phytochemicals enhance antioxidant enzyme expression to protect against NSAID-induced oxidative damage. *Molecular Nutrition & Food Research*, 61, 1600659.
29. Singh, D.P., Borse, S.P., Rana, R., & Nivsarkar, M. (2017). Curcumin efficiently prevents diclofenac sodium-induced gastroenteropathic damage in rats. *Food and Chemical Toxicology*, 108, 43–52.
30. Hatcher, H., Planalp, R., Cho, J., Torti, F.M., & Torti, S.V. (2008). Curcumin: From ancient medicine to current clinical trials. *Cellular and Molecular Life Sciences*, 65, 1631–1652.
31. Marchiani, A., Rozzo, C., Fadda, A., Delogu, G., & Ruzza, P. (2014). Curcumin and curcumin-like molecules: From spice to drugs. *Current Medicinal Chemistry*, 21, 204–222.
32. Hewlings, S.J., & Kalman, D.S. (2017). Curcumin: A review of its effects on human health. *Foods*, 6, 92.
33. Moghadamtousi, S.Z., Kadir, H.A., Hassandarvish, P., Tajik, H., Abubakar, S., & Zandi, K. (2014). A review on antibacterial, antiviral, and antifungal activity of curcumin. *BioMed Research International*, 2014, 186864.
34. Praditya, D., Kirchhoff, L., Brüning, J., Rachmawati, H., Steinmann, J., & Steinmann, E. (2019). Anti-infective properties of the golden spice curcumin. *Frontiers in Microbiology*, 10, 912.
35. Rai, M., Ingle, A.P., Pandit, R., Paralakar, P., Anasane, N., & Santos, C.A.D. (2020). Curcumin and curcumin-loaded nanoparticles: Antipathogenic and antiparasitic activities. *Expert Review of Anti-infective Therapy*, 18, 367–379.
36. Gupta, S.C., Patchva, S., & Aggarwal, B.B. (2013). Therapeutic roles of curcumin: Lessons learned from clinical trials. *AAPS Journal*, 15, 195–218.